

POPULATION:

All inpatients 18 years of age and older intubated and receiving partial mechanical ventilation support

SKILL LEVEL:

Licensed Respiratory Therapist

GENERAL CONSIDERATIONS:

- The Spontaneous Breathing Trial (SBT) will be performed with a low level CPAP (5 8 cm H₂O) while maintaining the ventilator's FiO₂
- Extubation Criteria:

A patient should be considered for extubation if all of the following are met:

- \circ A RSBI (or f/V_{T)} less than 100 after a 30 120 minute SBT (CPAP or PSV 5 H₂O)
- PaO₂/FiO₂ greater than 180 and PEEP less than 5 (CPAP) and FiO₂ less than 0.4
- Adequate airway protection, a reliable respiratory drive, and airway suctioning no more than every two hours
- o Successful "cuff leak test" in patients suspected of possible upper airway abnormalities
- o Arterial Blood Gases (ABG) are recommended 30 minutes post extubation

ORDERS:

- 1. Conduct a daily spontaneous breathing trial per Adult Mechanical Ventilation policy. Request minimum sedation before proceeding.
- 2. Notify provider and obtain order for extubation when Extubation Criteria is met

Signature of Physi	cian		
Date	Time		
Rev. 5.21		2005076103 CF (<i>Tab-Orders</i>)	